

Equivalences of Comodule Categories for Coalgebras over Rings.

Khaled AL-Takhman

Department of Mathematics

Birzeit University

P. O. Box 14, Birzeit, Palestine

E-mail: takhman@birzeit.edu

Abstract

In this article we defined and studied quasi-finite comodules, the cohom functors for coalgebras over rings. linear functors between categories of comodules are also investigated and it is proved that good enough linear functors are nothing but a cotensor functor. Our main result of this work characterizes equivalences between comodule categories generalizing the Morita-Takeuchi theory to coalgebras over rings. Morita-Takeuchi contexts in our setting is defined and investigated, a correspondence between strict Morita-Takeuchi contexts and equivalences of comodule categories over the involved coalgebras is obtained. Finally we proved that for coalgebras over QF-rings Takeuchi's representation of the cohom-functor is also valid.

Keywords: cotensor product, cohom functor, faithfully coflat comodules, equivalent categories, Morita-Takeuchi context.

Mathematics subject classification: 16W30

Introduction

M. Takeuchi [14] developed a theorem that characterizes equivalences of comodule categories over fields, dualizing Morita results on equivalences of module categories. In recent years a new interest arose for the study of coalgebras over rings. In this article we did this for equivalences of comodule categories.

One relevant consequence of the Morita theory on equivalences between module categories is that the notion of Morita equivalent rings is independent

categories of right modules are). Here, we extend the aforementioned equivalence theory to coalgebras over an arbitrary commutative ring R , under the mild hypothesis that the involved coalgebras are flat as R -modules. To transfer the Morita-Takeuchi theory from fields to commutative rings, we overcome two difficulties: on the one hand, no basis-dependent arguments can be used here, and, on the other, the lack of associativity in general of the cotensor product.

The cotensor functor plays the most important role in characterizing equivalences of comodule categories. S. Caenepeel [4] gave some sufficient conditions to guarantee the associativity of the cotensor product over rings, but unfortunately his result is not true (for a counter example see [6]). A detailed study of the cotensor functor for coalgebras over rings is done by the author in [2]. In section 2 we state some results on the exactness and associativity of this functor that we need in this work.

To prove our results we developed purely categorical arguments for the study of quasi-finite comodules and the cohom-functor in section 3. In this section we considered the coendomorphism coalgebra derived from a quasi-finite comodule X and showed that the dual algebra of this coalgebra is algebra (anti-) isomorphic to the algebra of comodule endomorphisms of X . In section 4 we studied linear functors (in particular equivalences) between comodule categories and showed that a linear functor, under certain conditions, is isomorphic to a cotensor functor. In section 5 we proved our main result (Theorem 5.5) which characterizes an equivalence by the existence of a bicomodule which is quasi-finite, faithfully coflat and an injector on either side.

The concept of Morita-Takeuchi context was first introduced in [14] for coalgebras over fields. In [5] it was studied for graded coalgebras. S. Caenepeel [4] defined it for coalgebras over rings, in his definition the purity condition which guarantees the associativity of the cotensor product is overlooked, so his definition for strict Morita-Takeuchi context does not give an equivalence between the comodule categories of the involved coalgebras. In section 6 we defined this concept for coalgebras over rings, and showed that there is a correspondence between strict Morita-Takeuchi contexts and equivalences of comodule categories. We also showed that for a C -comodule X that is quasi-finite, faithfully coflat and an injector the coendomorphism coalgebra of X is Morita-Takeuchi equivalent to C .

The ground ring has great influence on the properties of comodule categories. In section 7 we studied the case where the ring is a QF-ring and proved that most of the results of [14] are true for our settings.

Throughout this paper we assume that all rings are commutative with unity, all modules are unitary, and the unadorned tensor product is understood to be over the ground ring. The categorical terminology we used are those of

[12] with minor differences. For module theoretic notions we refer to [15]. Finally, a submodule W of an R -module V is called N -pure (N is an R -module) if the canonical map $W \otimes N \rightarrow V \otimes N$ is a monomorphism. W is called pure if it is N -pure for every N .

1 Notations and Preliminaries

Let R be a commutative ring with unity. We denote by $R\text{-Mod}$ the category of unital R -modules. In this section we recall the basic definitions and results that we need in the sequel.

Coalgebras : A *coalgebra* over a ring R is an R -module C together with two R -linear maps $\Delta : C \rightarrow C \otimes C$ (*comultiplication*) and $\varepsilon : C \rightarrow R$ (*counit*) such that $(id \otimes \Delta) \circ \Delta = (\Delta \otimes id) \circ \Delta$ (*Coassociativity property*) and $(\varepsilon \otimes id) \circ \Delta = id = (id \otimes \varepsilon) \circ \Delta$ (*counitary property*).

Coalgebra Morphisms : Let $\Delta_C : C \rightarrow C \otimes C$, $\Delta_D : D \rightarrow D \otimes D$ be coalgebras, an R -linear map $\pi : C \rightarrow D$ is called a *coalgebra morphism* if $\Delta_D \circ \pi = (\pi \otimes \pi) \circ \Delta_C$ and $\varepsilon_D \circ \pi = \varepsilon_C$.

Comodules : Let $\Delta : C \rightarrow C \otimes C$ be a coalgebra, a right C -comodule is an R -module M with an R -linear map $\varrho_M : M \rightarrow M \otimes C$ such that the following diagrams are commutative

$$\begin{array}{ccc} M & \xrightarrow{\varrho_M} & M \otimes C \\ \varrho_M \downarrow & & \downarrow id \otimes \Delta \\ M \otimes C & \xrightarrow{\varrho_M \otimes id} & M \otimes C \otimes C \end{array} \quad \begin{array}{ccc} M & \xrightarrow{\varrho_M} & M \otimes C \\ & \searrow \cong & \downarrow id \otimes \varepsilon \\ & & M \otimes R, \end{array}$$

which means, $(id \otimes \Delta) \circ \varrho_M = (\varrho_M \otimes id) \circ \varrho_M$ and $(id \otimes \varepsilon) \circ \varrho_M = id$. Left C -comodules are defined in a symmetric way.

Comodule Morphisms : Let $\varrho_M : M \rightarrow M \otimes C$, $\varrho_N : N \rightarrow N \otimes C$ be right C -comodules. An R -linear map $f : M \rightarrow N$ is called a *comodule morphism* (or C -colinear) if $\varrho_N \circ f = (f \otimes id) \circ \varrho_M$. The set of all C -comodule morphisms between M and N is denoted by $\text{Com}_C(M, N)$.

Example 1.1. Let M be a right C -comodule, for every R -module W , $W \otimes M$ has a structure of right C -comodule through $W \otimes M \xrightarrow{id \otimes \varrho_M} W \otimes M \otimes C$. With this structure on $M \otimes C$, ϱ_M becomes a right C -comodule morphism, which splits in $R\text{-Mod}$. Moreover, for each R -linear map $f : W \rightarrow V$ the map $f \otimes id : W \otimes M \rightarrow V \otimes M$ is C -colinear. The same can be done analogously for left C -comodules.

Bicomodules : Let C, D be two coalgebras, A C - D -bicomodule is a left C -comodule and a right D -comodule M , such that the left C -comodule structure map $\varrho_M^- : M \rightarrow C \otimes M$ is D -colinear, or equivalently the right D -comodule structure map $\varrho_M^+ : M \rightarrow M \otimes D$ is C -colinear.

Categories of Comodules : Let C and D be coalgebras. The right C -comodules together with the C -colinear maps between them constitute an additive category denoted by $\mathcal{M}^C$. Analogously, the categories of left C -comodules ${}^C\mathcal{M}$ and of C - D -bicomodules ${}^C\mathcal{M}^D$ can be defined.

Lemma 1.2. *Let C be a coalgebra. Then*

- (1) *The functor $- \otimes C : R\text{-Mod} \rightarrow \mathcal{M}^C$ is right adjoint to the forgetful functor by the natural isomorphisms (for $M \in \mathcal{M}^C, X \in R\text{-Mod}$),*

$$\text{Com}_C(M, X \otimes C) \rightarrow \text{Hom}_R(M, X), \quad f \mapsto (id \otimes \varepsilon) \circ f.$$

- (2) *For any $M \in \mathcal{M}^C$, the functor $- \otimes M : R\text{-Mod} \rightarrow \mathcal{M}^C$ is left adjoint to the functor $\text{Com}_C(M, -) : \mathcal{M}^C \rightarrow R\text{-Mod}$ by the natural isomorphisms (for $N \in \mathcal{M}^C, W \in R\text{-Mod}$),*

$$\text{Com}_C(W \otimes M, N) \rightarrow \text{Hom}_R(W, \text{Com}_C(M, N)), \quad f \mapsto [w \mapsto f \circ (w \otimes -)].$$

Proof: See [16, 6.11] ■

If C is flat as R -module, then the category $\mathcal{M}^C$ is a Grothendieck category (see [16]). Moreover a sequence

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

of C -comodules is exact in $\mathcal{M}^C$ iff it is exact in $R\text{-Mod}$, see [11].

From now on we assume that all coalgebras considered in this work are flat as R -modules.

2 The Cotensor Functor

The cotensor functor was first introduced by Milnor and More [9] for coalgebras over fields. Guzman [7, 8] studied this functor over rings but in the case of coseparable coalgebras. One of the key points to recover the Morita-Takeuchi theorem (see [14]) in our settings is the associativity of the cotensor product. In this section we state some results taken from [2] concerning the exactness and associativity of the cotensor functor.

Definition 2.1. Let $\varrho_M : M \rightarrow M \otimes C$ be a right and $\varrho_N : N \rightarrow C \otimes N$ a left C -comodule. The cotensor product of M and N (denoted $M \square_C N$) is defined as the kernel of the R -linear map

$$\alpha := \varrho_M \otimes id_N - id_M \otimes \varrho_N : M \otimes N \longrightarrow M \otimes C \otimes N.$$

So, we have the following exact sequence of R -modules

$$0 \rightarrow M \square_C N \rightarrow M \otimes N \xrightarrow{\alpha} M \otimes C \otimes N$$

For a right C -comodule M , the cotensor functor $M \square_C - : {}^C\mathcal{M} \rightarrow R\text{-Mod}$ is in general neither left nor right exact (see [1]), but we have the following

Lemma 2.2. *Let M be a right C -comodule, then*

- (1) *The functor $M \square_C - : {}^C\mathcal{M} \rightarrow R\text{-Mod}$ is (C, R) -left exact (also called relative left exact) i.e. left exact with respect to exact sequences of left C -comodules, that are pure in $R\text{-Mod}$.*
- (2) *If M is flat in $R\text{-Mod}$, then $M \square_C -$ is left exact.*

Tensor–Cotensor relation

Lemma 2.3. *Let $M \in \mathcal{M}^C$, $N \in {}^C\mathcal{M}$. For every $W \in R\text{-Mod}$ there exist two canonical R -linear maps*

$$\begin{aligned}\gamma_W &: W \otimes (M \square_C N) \longrightarrow (W \otimes M) \square_C N, \\ \mu_W &: (M \square_C N) \otimes W \longrightarrow M \square_C (N \otimes W).\end{aligned}$$

Moreover, the following are equivalent

- (1) $M \square_C N$ is W -pure in $M \otimes N$.
- (2) γ_W is an isomorphism.
- (3) μ_W is an isomorphism.

Remark: $W \otimes M$ and $N \otimes W$ have the trivial comodule structure given in example 1.1.

Associativity of the cotensor product

One of the main drawbacks of the behaviour of the cotensor product over rings is that it needs not be associative (see[6] for a counter example). But under some conditions it becomes so. Let M be a right C -comodule, L a C - D -bicomodule and N a left D -comodule. Then $M \square_C L$ has a structure of a right D -comodule through

$$M \square_C L \xrightarrow{id \square_C \varrho_L^+} M \square_C (L \otimes D) \cong (M \square_C L) \otimes D,$$

and it is a subcomodule of $M \otimes L$. Similarly $L \square_D N$ has a structure of left C -comodule.

Now we give a basic result from [2] that gives necessary conditions for the associativity of the cotensor product. For completeness the proof is included.

Proposition 2.4 (see [2]). *Let L be a C - D -bicomodule, M a right C -comodule and N a left D -comodule. If $M \square_C L$ is N -pure in $M \otimes L$ and $L \square_D N$ is M -pure in $L \otimes N$, then*

$$M \square_C (L \square_D N) \cong (M \square_C L) \square_D N.$$

Proof: $M \square_C L$ and $L \square_D N$ are right D - resp. left C -comodules, since C and D are flat in $R\text{-Mod}$. Consider the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & (M \square_C L) \square_D N & \longrightarrow & (M \otimes L) \square_D N & \longrightarrow & (M \otimes C \otimes L) \square_D N \\ & & \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 \\ 0 & \longrightarrow & M \square_C (L \square_D N) & \longrightarrow & M \otimes (L \square_D N) & \longrightarrow & M \otimes C \otimes (L \square_D N). \end{array}$$

The rows are exact: The first because $M \square_C L$ is N -pure in $M \otimes L$; the second because it defines $M \square_C (L \square_D N)$. Because of the M -purity of $L \square_D N$ in $L \otimes N$ and the flatness of C , ψ_2 and ψ_3 are isomorphisms. Hence ψ_1 is an isomorphism as required. $\blacksquare$

Coflat comodules

Definition 2.5. A right C -comodule M is called *coflat* (resp. *faithfully coflat*) if the functor $M \square_C - : {}^C\mathcal{M} \rightarrow R\text{-Mod}$ is exact (resp. exact and faithful).

Proposition 2.6. For a right C -comodule M the following are equivalent

1. M is faithfully coflat.
2. The functor $M \square_C - : {}^C\mathcal{M} \rightarrow R\text{-Mod}$ preserves and reflects exact sequences.
3. M is coflat and $M \square_C N \neq 0$ for every nonzero left C -comodule N .

Proof: This follows from general properties of functors between abelian categories. (See for example [12]) $\blacksquare$

If the ground ring is a field, then a C -comodule is coflat (resp. faithfully coflat) iff it is injective (resp. an injective cogenerator) (see [13]).

3 Quasi-finite comodules and the cohom functor

In this section we investigate the cohom functor, which was first introduced by M. Takeuchi [14] for coalgebras over fields. Some technical difficulties arise here, for example we lose the fact that over fields coflatness and injectivity are equivalent notions. To overcome this problem some injectivity preserving condition is imposed. We begin with the definition of quasi-finite comodules and the cohom functor. Some of the results given by Takeuchi concerning the cohom functor still hold in our general settings. In section 7 we recover all Takeuchi's results for coalgebras over QF-rings.

Definition 3.1. A right D -comodule X is called *quasi-finite*, if the functor $- \otimes X : R\text{-Mod} \rightarrow \mathcal{M}^D$ has a left adjoint. A quasi-finite left D -comodule is defined analogously.

Notation. 1. For a quasi-finite right D -comodule X , we denote the left adjoint of the functor $- \otimes X$ by $h_D(X, -) : \mathcal{M}^D \rightarrow R\text{-Mod}$. This functor is called the cohom functor. We also denote the adjunction bijection by

$$\Phi_{M,W} : \text{Hom}_R(h_D(X, M), W) \rightarrow \text{Com}_D(M, W \otimes X).$$

2. For every right D -comodule M we denote the unit of adjunction by $\eta_M : M \rightarrow h_D(X, M) \otimes X$, which satisfies that for every D -colinear map $f : M \rightarrow W \otimes X$, there exists a unique R -linear map $\tilde{f} : h_D(X, M) \rightarrow W$ such that $f = \Phi_{M,W}(\tilde{f}) = (\tilde{f} \otimes id)\eta_M$.

Remarks:. 1. Since the cohom functor has a right adjoint, it follows that it is right exact and commutes with direct limits.

2. From the definition it is clear that every quasi-finite comodule is flat in $R\text{-Mod}$.
3. The image of every $g \in \text{Hom}_R(h_D(X, M), W)$ under the adjunction bijection can be represented in terms of the counit by $\Phi_{M,W}(g) = (g \otimes id)\eta_M$.
4. For every right D -colinear map $f : N \rightarrow L$, there exists a unique R -linear map $h_D(X, f) : h_D(X, N) \rightarrow h_D(X, L)$ such that

$$\eta_L f = (h_D(X, f) \otimes id)\eta_N.$$

Some properties of the cohom functor

Let X be a quasi-finite right D -comodule, M a right D -comodule. From the above observations, for every R -module W , the D -colinear map

$$id \otimes \eta_M : W \otimes M \rightarrow W \otimes h_D(X, M) \otimes X$$

induces a unique canonical R -linear map

$$\lambda_W : h_D(X, W \otimes M) \rightarrow W \otimes h_D(X, M),$$

such that $(\lambda_W \otimes id_X)(\eta_{W \otimes M}) = id_W \otimes \eta_M$.

Proposition 3.2. *The map λ_W is an isomorphism.*

Proof: First it is easy to show that λ_W is an isomorphism for every free R -module W . Let W be an R -module and consider the free presentation of W , $R^{(\Lambda)} \rightarrow R^{(\Omega)} \rightarrow W \rightarrow 0$. By applying the functors $- \otimes M$, $h_D(X, -)$ and

$- \otimes h_D(X, M)$ to the free presentation of W we get the following commutative diagram with exact rows

$$\begin{array}{ccccccc}
R^{(\Lambda)} \otimes h_D(X, M) & \longrightarrow & R^{(\Omega)} \otimes h_D(X, M) & \longrightarrow & W \otimes h_D(X, M) & \longrightarrow & 0 \\
\uparrow \lambda_{R^{(\Lambda)}} & & \uparrow \lambda_{R^{(\Omega)}} & & \uparrow \lambda_W & & \\
h_D(X, R^{(\Lambda)} \otimes M) & \longrightarrow & h_D(X, R^{(\Omega)} \otimes M) & \longrightarrow & h_D(X, W \otimes M) & \longrightarrow & 0,
\end{array}$$

in which $\lambda_{R^{(\Omega)}}$ and $\lambda_{R^{(\Lambda)}}$ are isomorphisms, hence λ_W is an isomorphism. $\blacksquare$

Corollary 3.3. Let X be a quasi-finite right D -comodule, M a right D -comodule, then we have the following isomorphism of functors,

$$h_D(X, - \otimes M) \cong - \otimes h_D(X, M) : R\text{-Mod} \rightarrow R\text{-Mod}.$$

Proof: For all $V, W \in R\text{-Mod}$ we have the natural isomorphisms

$$\begin{aligned}
\text{Hom}_R(h_D(X, V \otimes M), W) &\cong \text{Com}_D(V \otimes M, W \otimes X) \\
&\cong \text{Hom}_R(V, \text{Com}_D(M, W \otimes X)) \\
&\cong \text{Hom}_R(V, \text{Hom}_R(h_D(X, M), W)) \\
&\cong \text{Hom}_R(V \otimes h_D(X, M), W)
\end{aligned}$$

$\blacksquare$

Comodule Structure on $h_D(X, M)$

In the following we consider the behaviour of the cohom functor with respect to bicomodule structures on its arguments. It follows that the cohom functor is a bifunctor $h(-, -) : {}^C\mathcal{M}^D \times {}^E\mathcal{M}^D \rightarrow {}^E\mathcal{M}^C$ contravariant in the first and covariant in the second.

Lemma 3.4. (1) Let X be a quasi-finite right D -comodule, M a C - D -bicomodule, then the R -linear map

$$h_D(X, \varrho_M^-) : h_D(X, M) \rightarrow h_D(X, C \otimes M) \cong C \otimes h_D(X, M),$$

gives $h_D(X, M)$ a structure of a left C -comodule. With this structure the map $\eta_M : M \rightarrow h_D(X, M) \otimes X$ is C - D -bilinear.

(2) Let X be a C - D -bicomodule, M a right D -comodule. If X_D is quasi-finite, then there exists an R -linear map

$$\varrho_h : h_D(X, M) \longrightarrow h_D(X, M) \otimes C$$

which gives $h_D(X, M)$ a structure of right C -comodule, therefore we have a functor $h_D(X, -) : \mathcal{M}^D \rightarrow \mathcal{M}^C$. Moreover we have $\text{Im}(\eta_M) \subseteq h_D(X, M) \square_C X$, where η_M is the unit of adjunction.

Proof: (1) Follows directly from the properties of the unit of adjunction.

- (2) The D -colinear map $M \xrightarrow{(id \otimes \varrho_X^-) \eta_M} h_D(X, M) \otimes C \otimes X$, where $\varrho_X^- : X \rightarrow C \otimes X$ is the left C -comodule structure map, induces the unique R -linear map

$$\varrho_h : h_D(X, M) \rightarrow h_D(X, M) \otimes C,$$

such that $(id \otimes \varrho_X^-) \eta_M = (\varrho_h \otimes id) \eta_M$. This map gives $h_D(X, M)$ a structure of a right C -comodule. To show that $h_D(X, -) : \mathcal{M}^D \rightarrow \mathcal{M}^C$ defines a functor, we have to show that, for every right D -colinear map $f : M \rightarrow N$, the induced map $h_D(X, f) : h_D(X, M) \rightarrow h_D(X, N)$ is C -colinear. This follows from the uniqueness of the R -linear map which is induced by the D -colinear map $M \xrightarrow{(id \otimes \varrho_X^-) \circ \eta_N \circ f} h_D(X, N) \otimes C \otimes X$.

For the other assertion we have

$$\begin{aligned} (\varrho_h \otimes id - id \otimes \varrho_X^-) \eta_M &= (\varrho_h \otimes id) \eta_M - (id \otimes \varrho_X^-) \eta_M \\ &= (\varrho_h \otimes id) \eta_M - (\varrho_h \otimes id) \eta_M \\ &= 0, \end{aligned}$$

hence, $\text{Im}(\eta_M) \subseteq \text{Ker}(\varrho_h \otimes id - id \otimes \varrho_X^-) = h_D(X, M) \square_C X$. ■

The following corollary follows directly from the above constructions.

Corollary 3.5. Let X be a quasi-finite right D -comodule.

- (1) For every right D -comodule M there exists a unique R -linear map

$$\delta_M : h_D(X, M) \rightarrow M \square_D h_D(X, D), \quad \text{with } (\delta_M \otimes id) \eta_M = id \square_D \eta_D$$

- (2) If X is also a C - D -bicomodule, then δ_M is right C -colinear. If M is a C - D -bicomodule, then δ_M is C -bilinear

The following theorem was given by Takeuchi [14] for coalgebras over fields, and it holds also for coalgebras over rings.

Theorem 3.6. Let X be a C - D -bicomodule, then the following are equivalent

1. X_D is quasi-finite.
2. The functor $-\square_C X : \mathcal{M}^C \rightarrow \mathcal{M}^D$ has a left adjoint. In this case the left adjoint of $-\square_C X$ is the functor $h_D(X, -) : \mathcal{M}^D \rightarrow \mathcal{M}^C$.

Proof: (1) $\Rightarrow$ (2). Define

$$\begin{aligned} \text{Com}_C(h_D(X, M), N) &\rightleftharpoons \text{Com}_D(M, N \square_C X) \\ f &\longmapsto (f \square_C id) \eta_M \\ \tilde{g} &\longleftarrow g, \end{aligned}$$

where $\tilde{g}$ is the unique R -linear map $\tilde{g} : h_D(X, M) \rightarrow N$ such that $g = (\tilde{g} \otimes id) \eta_X$. It is straightforward to see that $(f \square_C id) \eta_M$ is D -colinear, and that the two maps are inverse to each other. So we have only to show that $\tilde{g}$ is C -colinear. We have

$$\begin{aligned} (\varrho_N \tilde{g} \otimes id) \eta_X &= (\varrho_N \otimes id)(\tilde{g} \otimes id) \eta_X \\ &= (\varrho_N \otimes id) g. \end{aligned}$$

and

$$\begin{aligned} ((\tilde{g} \otimes id) \varrho_h \otimes id) \eta_X &= (\tilde{g} \otimes id \otimes id)(\varrho_h \otimes id) \eta_X \\ &= (\tilde{g} \otimes id \otimes id)(id \otimes \varrho_X^-) \eta_X \\ &= (\tilde{g} \otimes \varrho_X^-) \eta_X \\ &= (id \otimes \varrho_X^-)(\tilde{g} \otimes id) \eta_X \\ &= (id \otimes \varrho_X^-) g. \end{aligned}$$

Now $(\varrho_N \otimes id - id \otimes \varrho_X^-)|_{N \square_C X} = 0$, hence $((\tilde{g} \otimes id) \varrho_h \otimes id) \eta_X = (\varrho_N \tilde{g} \otimes id) \eta_X$. But an R -linear map $\mu : h_D(X, M) \rightarrow N \otimes C$, such that $(id \otimes \varrho_X^-)g = (\mu \otimes id) \eta_X$ is unique, hence $(\tilde{g} \otimes id) \varrho_h = \varrho_N \tilde{g}$, i.e $\tilde{g}$ is C -colinear.

(2) $\Rightarrow$ (1). Assume that $-\square_C X$ has a left adjoint. The functor $-\otimes X : R\text{-Mod} \rightarrow \mathcal{M}^D$ can be written as the composition of the functors $-\otimes C : R\text{-Mod} \rightarrow \mathcal{M}^C$, which has a left adjoint (see 1.2), and $-\square_C X : \mathcal{M}^C \rightarrow R\text{-Mod}$, this is because for all $W \in R\text{-Mod}$, we have $(W \otimes C) \square_C X \cong W \otimes X$. $\blacksquare$

Corollary 3.7. Let X be a C - D -bicomodule, X_D quasi-finite. For every D -comodule M there exists a D -colinear map (the unit of adjunction) $\eta_M : M \rightarrow h_D(X, M) \square_C X$, such that, for every D -colinear map $f : M \rightarrow N \square_C X$ ($N \in \mathcal{M}^C$), there exists a unique C -colinear map $\tilde{f} : h_D(X, M) \rightarrow N$ with $f = (\tilde{f} \square_C id) \eta_M$. Moreover, if M is a C - D -bicomodule, then η_M is C - D -bilinear.

Exactness of the cohom functor

Now we consider the questions: Under what conditions is the cohom functor exact; and what happens if this is the case.

Proposition 3.8. Let X be a quasi-finite right D -comodule. The following are equivalent

- (1) The cohom functor $h_D(X, -) : \mathcal{M}^D \rightarrow R\text{-Mod}$ is exact;

(2) $W \otimes X$ is injective in $\mathcal{M}^D$, for every injective R -module W .

If X is a C - D -bicomodule, then the following are also equivalent to (1) and (2) above.

(3) The cohom functor $h_D(X, -) : \mathcal{M}^D \rightarrow \mathcal{M}^C$ is exact;

(4) $N \square_C X$ is injective in $\mathcal{M}^D$, for every injective right C -comodule N .

Proof: This is clear, since $h_D(X, -)$ is left adjoint to $- \otimes X$ and the category $\mathcal{M}^D$ has enough injectives. ■

Analogous to the definition of injectors in module categories (see [3, Exercise 21.7]), we define injectors in comodule categories.

Definition 3.9. Let M be a right C -comodule. M is said to be an injector in $\mathcal{M}^C$ if the functor $- \otimes M : R\text{-Mod} \rightarrow \mathcal{M}^C$ respects injective objects. Injectors in ${}^C\mathcal{M}$ are defined similarly. C itself as a comodule is an injector in $\mathcal{M}^C$ and in ${}^C\mathcal{M}$.

Corollary 3.10. Let X be a C - D -bicomodule with X_D quasi-finite, then the following are equivalent

- (a) X_D is an injector
- (b) the cohom functor $h_D(X, -) : \mathcal{M}^D \rightarrow \mathcal{M}^C$ is exact
- (c) the functor $- \square_C X : \mathcal{M}^C \rightarrow \mathcal{M}^D$ respects injective objects.

Now we show that the cohom functor under certain condition is nothing but a cotensor functor.

Proposition 3.11. Let X be a quasi-finite D -comodule. If the cohom functor is exact, then

- (1) $h_D(X, M)$ is flat in $R\text{-Mod}$, for every D -comodule M , which is flat in $R\text{-Mod}$,
- (2) the R -linear map δ_M in 3.5 is an isomorphism, for every right C -comodule M ,
- (3) if, in addition, X is a C - D -bicomodule, then the δ is a bijective C -colinear map. If M is also a C - D -bicomodule, then δ is a bijective C -bilinear map.

Proof: (1) We know that $- \otimes h_D(X, M) \cong h_D(X, - \otimes M)$.

(2) let $M \in \mathcal{M}^D$. The sequence

$$0 \rightarrow M \xrightarrow{\varrho_M} M \otimes D \xrightarrow{\varrho_M \otimes id - id \otimes \Delta} M \otimes D \otimes D$$

is exact in $\mathcal{M}^D$. The result follows now from the following diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & h_D(X, M) & \longrightarrow & h_D(X, M \otimes D) & \longrightarrow & h_D(X, M \otimes D \otimes D) \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & M \square_D h_D(X, D) & \longrightarrow & M \otimes h_D(X, D) & \longrightarrow & M \otimes D \otimes h_D(X, D). \end{array}$$

(3) Follows directly from 2. ■

Corollary 3.12. Let X be a quasi-finite D -comodule. If the cohom functor is exact then we have

$$h_D(X, -) \cong -\square_D h_D(X, D) : \mathcal{M}^D \rightarrow R\text{-Mod}$$

Proof: The isomorphism δ_M in 3.11 is natural. ■

The coendomorphism coalgebra

For a quasi-finite D -comodule, and dual to the notion of endomorphism algebras for modules, the R -module $h_D(X, X)$ has a structure of a coalgebra. This structure was considered in [14] for fields, and it is also valid in our case.

Let X be a quasi-finite D -comodule, let $e_D(X) := h_D(X, X)$. This R -module has a structure of an R -coalgebra, where the comultiplication corresponds to the D -colinear map

$$(id \otimes \eta_X)\eta_X : X \rightarrow h_D(X, X) \otimes h_D(X, X) \otimes X$$

in the adjunction bijection for $M = X$. The counit corresponds to the D -colinear map $id : X \rightarrow R \otimes X$ for $M = X$ and $W = R$. This R -coalgebra is called the *coendomorphism coalgebra* of X . The unit of adjunction $\eta_X : X \rightarrow h_D(X, X) \otimes X$ gives X a structure of a left $e_D(X)$ -comodule, hence X is a $e_D(X)$ - D -coalgebra.

It is well known that for an R -coalgebra $\Delta : C \rightarrow C \otimes C$, the dual R -module $C^* = \text{Hom}_R(C, R)$ is an R -algebra under the convolution product (i.e. for $f, g \in C^*$, $f * g = (f \otimes g)\Delta$). We now give a result about the dual algebra of the coendomorphism coalgebra.

Proposition 3.13. *For a quasi-finite right D -comodule X , the dual algebra of the coendomorphism coalgebra is algebra anti-isomorphic to the algebra of right D -comodule endomorphisms of X . (i.e. $(e_D(X))^* \cong \text{Com}_D(X, X)$).*

Proof: Let X be a quasi-finite right D -comodule, and consider the composition of the maps

$$e_D(X)^* = \text{Hom}_R(h_D(X, X), R) \xrightarrow[\cong]{\Phi_{X,R}} \text{Com}_D(X, R \otimes X) \cong \text{Com}_D(X, X).$$

The image of $f \in e_D(X)^*$ under this composition is $\vartheta(f \otimes id) \circ \eta_X$, where $\vartheta : R \otimes X \rightarrow X$ is the canonical isomorphism. It is easy to see that this map is an algebra anti-isomorphism. $\blacksquare$

Remark:. In 3.13, if we start with a left quasi-finite D -comodule, then the algebra anti-isomorphism there is an algebra isomorphism.

4 Functors between comodule categories

Let C, D be R -coalgebras, we will study additive functors $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$, which are assumed to be R -linear, i.e the canonical maps $\text{Com}_C(M, N) \rightarrow \text{Com}_D(F(M), F(N))$, induced by F for $M, N \in \mathcal{M}^C$, are assumed to be R -linear. We will show that good enough R -linear functors are isomorphic to a cotensor functor. Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be an R -linear functor. We consider the bifunctors

$$- \otimes F(-), F(- \otimes -) : R\text{-Mod} \times \mathcal{M}^C \rightarrow \mathcal{M}^D,$$

where for $W \in R\text{-Mod}$, $M \in \mathcal{M}^C$, $W \otimes M$ (resp. $W \otimes F(M)$) is endowed with the canonical structure of right C -comodule (resp. D -comodule). Our aim is to construct a natural transformation

$$\lambda : - \otimes F(-) \rightarrow F(- \otimes -).$$

For every C -comodule M let $\bar{\lambda}_{R,M}$ be the unique isomorphism, that render the following diagram commutative

$$\begin{array}{ccc} R \otimes F(M) & \xrightarrow{\bar{\lambda}_{R,M}} & F(R \otimes M) \\ \cong \downarrow & & \downarrow \cong \\ F(M) & \xrightarrow{=} & F(M), \end{array}$$

where $M \in \mathcal{M}^C$ and the vertical arrows are the canonical isomorphisms.

Lemma 4.1. *Let $\mathcal{P} = \{R\}$ be the full subcategory of $R\text{-Mod}$ whose only object is R , denote the restriction of the functors $- \otimes F(M)$ and $F(- \otimes M)$ on $\mathcal{P}$ by T and S resp. Then $\bar{\lambda}_{-,M} : T \rightarrow S$ is a natural transformation.*

Proof: Given a homomorphism $f : R \rightarrow R$ in $\mathcal{P}$, we have to check that the diagram

$$\begin{array}{ccc} R \otimes F(M) & \xrightarrow{f \otimes id} & R \otimes F(M) \\ \bar{\lambda}_{R,M} \downarrow & & \downarrow \bar{\lambda}_{R,M} \\ F(R \otimes M) & \xrightarrow{F(f \otimes id)} & F(R \otimes M) \end{array}$$

commutes.

Define $g : F(M) \rightarrow F(M)$, by $g(x) = f(1)x$. It is easy to see that g is D -colinear and that $g = F(\tilde{g})$, where $\tilde{g} : M \rightarrow M, m \mapsto f(1)m$.

Consider the diagram

$$\begin{array}{ccccc} & & F(M) & \xrightarrow{g} & F(M) \\ & \nearrow \cong & \downarrow & & \downarrow id \\ R \otimes F(M) & \xrightarrow{f \otimes id} & R \otimes F(M) & & \\ \bar{\lambda}_{R,M} \downarrow & & \downarrow id & & \downarrow \bar{\lambda}_{R,M} \\ & \nearrow \cong & F(M) & \xrightarrow{g} & F(M) \\ F(R \otimes M) & \xrightarrow{F(f \otimes id)} & F(R \otimes M) & & \end{array}$$

By the definition of $\bar{\lambda}_{R,M}$ the right and left rectangles are commutative, the upper rectangle is commutative because $f(r) = f(1)r$ for all $r \in R$. The lower rectangle is commutative, since it is obtained from the diagram

$$\begin{array}{ccc} M & \xrightarrow{\tilde{g}} & M \\ \cong \uparrow & & \uparrow \cong \\ R \otimes M & \xrightarrow{f \otimes id} & R \otimes M \end{array}$$

which is commutative. Now, since the map $F(R \otimes M) \rightarrow F(M)$ is a monomorph (isomorph), the front rectangle is commutative. This implies that $\bar{\lambda}_{-,M} : - \otimes F(M) \rightarrow F(- \otimes M)$ is natural, where $- \otimes F(M), F(- \otimes M) : \mathcal{P} \rightarrow \mathcal{M}^D$. ■

By a theorem of Mitchel ([10, Theorem 3.6.5]), $\bar{\lambda}_{-,M}$ is uniquely extended to a natural transformation

$$\lambda_{-,M} : - \otimes F(M) \rightarrow F(- \otimes M),$$

for every right C -comodule M , where now $- \otimes F(M), F(- \otimes M) : R\text{-Mod} \rightarrow \mathcal{M}^D$. Moreover (see [10, Corollary 3.6.6]), if F preserves direct sums (resp. direct limits, inductive limits), then $\lambda_{W,-}$ is a natural isomorphism for every projective R -module (resp. flat R -module, R -module) W . Now we want to see under what conditions is $\lambda_{W,M}$ functorial in M .

Lemma 4.2. Assume that $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ respects direct sums, then

$$\lambda_{-, -} : - \otimes F(-) \rightarrow F(- \otimes -)$$

is a natural transformation. Moreover, if F preserves direct limits, then $\lambda_{W, -}$ is a natural isomorphism for every flat R -module W . Finally, if F preserves inductive limits, then $\lambda_{-, -}$ is a natural isomorphism.

Proof: We know that $\lambda_{W, M}$ is functorial in W . Thus we have to show that it is functorial in M . Let $g : M \rightarrow N$ be C -colinear. From the commutative diagram

$$\begin{array}{ccccc} R \otimes F(M) & \xrightarrow{id \otimes F(f)} & R \otimes F(N) & & \\ \downarrow \lambda_{R, M} & \searrow \cong & \downarrow & \searrow \cong & \\ & F(M) & \xrightarrow{F(f)} & F(N) & \\ & \uparrow \cong & \downarrow \lambda_{R, N} & \uparrow \cong & \\ F(R \otimes M) & \xrightarrow{F(id \otimes f)} & F(R \otimes N) & & \end{array}$$

we obtain that $\lambda_{R, M}$ is functorial in M . Now, since F preserves direct sums, $\lambda_{W, M}$ is natural for every free R -module W . In the general case use a free presentation for W to show that $\lambda_{W, M}$ is natural (in M) for every R -module W . The other assertions follow from the above observations. $\blacksquare$

Bicomodule structure on $F(C)$

Now, using the natural transformation λ and by imposing a mild condition on F , we want to give $F(C)$ a structure of a bicomodule. To this end we need the following lemma

Lemma 4.3. Assume that $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ preserves direct sums, then for every $W \in R\text{-Mod}$ and $M, N \in \mathcal{M}^C$, the following formula holds

$$\lambda_{W, M \otimes N} \circ (id \otimes \lambda_{M, N}) = \lambda_{W \otimes M, N}$$

Proof: It is easy to prove the result for $W = R$ and hence for every free R -module W . Now let $W \in R\text{-Mod}$, from the free presentation $R^{(\Lambda)} \xrightarrow{\theta} W \rightarrow 0$ we obtain the following diagram

$$\begin{array}{ccccc} R^{(\Lambda)} \otimes F(M \otimes N) & \xrightarrow{\theta \otimes F(id)} & W \otimes F(M \otimes N) & & \\ \downarrow \lambda_{R^{(\Lambda)}, M \otimes N} & \swarrow id \otimes \lambda_{M, N} & \downarrow & \swarrow id \otimes \lambda_{M, N} & \\ & R^{(\Lambda)} \otimes M \otimes F(N) & \xrightarrow{\theta \otimes id} & W \otimes M \otimes F(N) & \\ & \swarrow \lambda_{R^{(\Lambda)} \otimes M, N} & \downarrow \lambda_{W, M \otimes N} & \swarrow \lambda_{W \otimes M, N} & \\ F(R^{(\Lambda)} \otimes M \otimes N) & \xrightarrow{F(\theta \otimes id)} & F(W \otimes M \otimes N) & & \end{array}$$

in which all sub-diagrams (except possibly the right triangle) are commutative and $\theta \otimes id$ is an epimorph. Hence the right triangle is commutative and the result follows. $\blacksquare$

Proposition 4.4. *Assume that F preserves direct limits, then $F(C)$ has a structure of a left C -comodule. Hence $F(C)$ becomes a C - D -bicomodule.*

Proof: Since F preserves direct limits, $\lambda_{C,C}$ is an isomorphism, since C_R is flat. Let $\varrho_{F(C)} : F(C) \longrightarrow C \otimes F(C)$ be the unique R -linear map that makes the diagram

$$\begin{array}{ccc} F(C) & \xrightarrow{\varrho_{F(C)}} & C \otimes F(C) \\ & \searrow F(\Delta) & \swarrow \lambda_{C,C} \\ & F(C \otimes C) & \end{array}$$

commutative.

We will show that $\varrho_{F(C)}$ is a comodule structure map. First, we have to show that the following diagram is commutative

$$\begin{array}{ccc} F(C) & \xrightarrow{\varrho_{F(C)}} & C \otimes F(C) \\ \varrho_{F(C)} \downarrow & (*) & \downarrow \Delta \otimes id \\ C \otimes F(C) & \xrightarrow{id \otimes \varrho_{F(C)}} & C \otimes C \otimes F(C). \end{array}$$

Imbed this diagram in the following diagram

$$\begin{array}{ccccc} & & F(C) & \xrightarrow{\varrho_{F(C)}} & C \otimes F(C) \\ & \swarrow \varrho_{F(C)} & \downarrow & \searrow \Delta \otimes id & \downarrow \lambda_{C,C} \\ C \otimes F(C) & \xrightarrow{id \otimes \varrho_{F(C)}} & C \otimes C \otimes F(C) & & \\ \downarrow \lambda_{C,C} & \swarrow id \otimes F(\Delta) & \downarrow id & \swarrow id \otimes \lambda_{C,C} & \downarrow \lambda_{C \otimes C, C} \\ & C \otimes F(C \otimes C) & & & \\ & \downarrow & & & \\ & F(C) & \xrightarrow{F(\Delta)} & F(C \otimes C) & \\ \swarrow F(\Delta) & & \downarrow \lambda_{C,C \otimes C} & \swarrow F(\Delta \otimes id) & \\ F(C \otimes C) & \xrightarrow{F(id \otimes \Delta)} & F(C \otimes C \otimes C) & & \end{array}$$

This diagram (except possibly the top rectangle) is commutative, this is shown using lemma 4.3, the coassociativity of Δ , that $\lambda_{C \otimes C, C}$ is an isomorphism, and that $\lambda_{-, C}$ is natural. Hence the top side is commutative since $\lambda_{C \otimes C, C}$ is a monomorphism.

To show that $(\varepsilon \otimes id)\varrho_{F(C)} = id$, consider the diagram

$$\begin{array}{ccccc}
 R \otimes F(C) & \xrightarrow{\lambda_{R, C}} & F(R \otimes C) & & \\
 \uparrow \varepsilon \otimes id & \nwarrow \cong & \nearrow \cong & & \uparrow F(\varepsilon \otimes id) \\
 & F(C) & & & \\
 & \nwarrow \varrho_{F(C)} & \nearrow F(\Delta) & & \\
 C \otimes F(C) & \xrightarrow{\lambda_{C, C}} & F(C \otimes C) & &
 \end{array}$$

which is shown to be commutative by the counitary property, and the definition of $\varrho_{F(C)}$. $\blacksquare$

We know that the cotensor functor is relative left exact and respects direct limits. Now we give a theorem, similar to Watts theorem for modules, which shows that a functor that is relative left exact and respects direct limits with an extra condition can be presented as a cotensor functor.

Theorem 4.5. *Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be a relative left exact functor that respects direct limits. If $\lambda_{M, C}$ and $\lambda_{M \otimes C, C}$ are isomorphisms, for every right C -comodule M (e.g. if F respects inductive limits or F is an equivalence), then F is naturally isomorphic to $-\square_C F(C)$.*

Proof: Let $\varrho_M : M \rightarrow M \otimes C$ be a right C -comodule. We have the following (C, R) -exact sequence of comodules

$$0 \longrightarrow M \longrightarrow M \otimes C \longrightarrow M \otimes C \otimes C,$$

hence we get the following commutative (see 4.3) diagram with exact rows

$$\begin{array}{ccccccc}
 0 \longrightarrow & M \square_C F(C) & \longrightarrow & M \otimes F(C) & \xrightarrow{\varrho_M \otimes id - id \otimes \varrho_{F(C)}} & M \otimes C \otimes F(C) & \\
 & \downarrow \phi_M & & \downarrow \lambda_{M, C} & & \downarrow \lambda_{M \otimes C, C} & \\
 0 \longrightarrow & F(M) & \longrightarrow & F(M \otimes C) & \xrightarrow{F(\varrho_M \otimes id - id \otimes \Delta)} & F(M \otimes C \otimes C), &
 \end{array}$$

where the desired isomorphism ϕ_M is given by the universal property of the kernel. To show the naturality of ϕ_M , let $f : M \rightarrow N$ be a C -colinear map

and use the following commutative diagram

$$\begin{array}{ccccc}
& & M \otimes F(C) & \xrightarrow{f \otimes id} & N \otimes F(C) \\
& \nearrow & \downarrow & & \nearrow \\
M \square_C F(C) & \xrightarrow{f \square_C id} & N \square_C F(C) & & \\
\downarrow \phi_M & & \downarrow \lambda_{M,C} & & \downarrow \lambda_{N,C} \\
& & F(M \otimes C) & \xrightarrow{F(f \otimes id)} & F(N \otimes C) \\
& \nearrow F(\varrho_M) & \downarrow \phi_N & & \nearrow F(\varrho_N) \\
F(M) & \xrightarrow{F(f)} & F(N) & &
\end{array}$$

to see that ϕ_M is functorial in M . ■

Corollary 4.6. Let C, D be R -coalgebras. If $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ is a category equivalence with inverse $G : \mathcal{M}^D \rightarrow \mathcal{M}^C$, then there exist bicomodules $X \in {}^C\mathcal{M}^D$ and $Y \in {}^D\mathcal{M}^C$ such that

$$F \cong - \square_C X \quad \text{and} \quad G \cong - \square_D Y.$$

Proof: $X = F(C)$ and $Y = G(D)$. ■

5 Equivalences of Comodule Categories

In this section we study equivalences between comodule categories, and under what conditions we get such an equivalence. We prove a generalization of Morita-Takeuchi theorem for our settings. We begin with some properties of the bicomodules $F(C)$ and $G(D)$ in corollary 4.6.

Lemma 5.1. Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be a category equivalence with inverse $G : \mathcal{M}^D \rightarrow \mathcal{M}^C$. Then the following hold

- (1) $F(C)$ and $G(D)$ are flat R -modules.
- (2) For each right C -comodule M , $M \square_C F(C)$ is a pure submodule of $M \otimes F(C)$.
- (3) For each right D -comodule N , $N \square_D G(D)$ is pure in $N \otimes G(D)$.
- (4) For every $P \in {}^D\mathcal{M}$ we have

$$(G(D) \square_C F(C)) \square_D P \cong G(D) \square_C (F(C) \square_D P)$$

- (5) For every $Q \in {}^C\mathcal{M}$ we have

$$(F(C) \square_D G(D)) \square_C Q \cong F(C) \square_D (G(D) \square_C Q).$$

Proof: (1) $- \otimes F(C) \cong F(- \otimes C)$ which is exact, since C is flat in $R\text{-Mod}$ and F is an equivalence. Similarly for $G(D)$.

(2) Let $M \in \mathcal{M}^C$. For every $W \in R\text{-Mod}$ we have the canonical R -linear map $\gamma_W : W \otimes (M \square_C F(C)) \rightarrow (W \otimes M) \square_C F(C)$, (see lemma 2.3). Consider the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & F(W \otimes M) & \longrightarrow & F(W \otimes M \otimes C) & \longrightarrow & F(W \otimes M \otimes C \otimes C) \\
& & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
0 & \longrightarrow & W \otimes (M \square_C F(C)) & \longrightarrow & W \otimes (M \otimes F(C)) & \longrightarrow & W \otimes (M \otimes C \otimes F(C)) \\
& & \downarrow \gamma_W & & \downarrow \cong & & \downarrow \cong \\
0 & \longrightarrow & (W \otimes M) \square_C F(C) & \longrightarrow & (W \otimes M) \otimes F(C) & \longrightarrow & (W \otimes M) \otimes C \otimes F(C),
\end{array}$$

in which the first and third rows are exact, hence the second is also exact. The result follows now from lemma 2.3.

(3) Similar to 2.

(4) The assertions in (4) and (5) follow from (1), (2) and proposition 2.4. ■

With the help of representing an equivalence by a cotensor functor and lemma 5.1, we can now prove that an equivalence between right comodule categories gives an equivalence of the corresponding left comodule categories

Theorem 5.2. *Let C, D be two coalgebras. If $\mathcal{M}^C$ is equivalent to $\mathcal{M}^D$, then ${}^C\mathcal{M}$ is equivalent to ${}^D\mathcal{M}$.*

Proof: Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be an equivalence with inverse $G : \mathcal{M}^D \rightarrow \mathcal{M}^C$. Define $F' := F(C) \square_D - : {}^D\mathcal{M} \rightarrow {}^C\mathcal{M}$ and $G' := G(D) \square_C - : {}^C\mathcal{M} \rightarrow {}^D\mathcal{M}$. For $P \in {}^D\mathcal{M}$, $Q \in {}^C\mathcal{M}$, we have

$$\begin{aligned}
F'G'(Q) &= F(C) \square_D (G(D) \square_C Q) \\
&\cong (F(C) \square_D G(D)) \square_C Q \\
&\cong GF(C) \square_C Q \\
&\cong C \square_C Q \cong Q. \\
G'F'(P) &= G(D) \square_C (F(C) \square_D P) \\
&\cong (G(D) \square_C F(C)) \square_D P \\
&\cong FG(D) \square_D P \\
&\cong D \square_D P \cong P.
\end{aligned}$$

Hence F' is an equivalence with inverse G' ■

Definition 5.3. Two coalgebras are called Morita-Takeuchi equivalent if the categories of right (equivalently of left) comodules over these coalgebras are equivalent

The bicomodules $F(C)$ and $G(D)$ have other properties

Corollary 5.4. Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be a category equivalence with inverse $G : \mathcal{M}^D \rightarrow \mathcal{M}^C$. Then $F(C)$ is quasi-finite, faithfully coflat and an injector as right D - and as left C -comodule. Moreover $e_D(F(C)) \cong C$ and $e_C(F(C)) \cong D$ as coalgebras. Similar results hold for $G(D)$.

Proof: The fact that $F \cong -\square_C F(C)$ is an equivalence with inverse $G \cong -\square_D G(D)$ implies that $F(C)$ is quasi-finite and an injector as right D -comodule and faithfully coflat as left C -comodule, and that $e_D(F(C)) \cong F(C) \square_D G(D) \cong GF(D) \cong C$. The other assertions follow from the fact that $F(C) \square_D -$ is an equivalence with inverse $G(D) \square_C -$. ■

We are now ready to give our main result in this article, which generalizes the Morita-Takeuchi theorem.

Theorem 5.5. *Let C and D be two coalgebras. The following are equivalent.*

- (1) *C and D are Morita-Takeuchi equivalent.*
- (2) *The categories ${}^C\mathcal{M}$ and ${}^D\mathcal{M}$ are equivalent.*
- (3) *There exists a C - D -bicomodule X , such that X_D is quasi-finite, faithfully coflat and an injector, and $e_D(X) \cong C$ as coalgebras.*
- (4) *There exists a C - D -bicomodule X , such that ${}_CX$ is quasi-finite, faithfully coflat and an injector, and $e_C(X) \cong D$ as coalgebras.*
- (5) *There exists a D - C -bicomodule Y , such that Y_C is quasi-finite, faithfully coflat and an injector, and $e_C(X) \cong D$ as coalgebras.*
- (6) *There exists a D - C -bicomodule Y , such that ${}_DY$ is quasi-finite, faithfully coflat and an injector, and $e_D(Y) \cong C$ as coalgebras.*

Proof: From what we have done, we have only to show that (3) $\Rightarrow$ (1). Assume that a C - D -bicomodule X satisfies the conditions of (3). We show that $-\square_C X : \mathcal{M}^C \rightarrow \mathcal{M}^D$ is an equivalence.

From corollary 3.12 we know that $h_D(X, -) \cong -\square_D h_D(X, D)$, hence $h_D(X, D)$ is a coflat D -comodule. Next we show that $\eta_D : D \rightarrow h_D(X, D) \square_C X$, (see 3.7), is an isomorphism. The C -bilinear map $\delta_X : h_D(X, X) \cong C \rightarrow X \square_D h_D(X, D)$ is an isomorphism (see 3.11) and the diagram

$$\begin{array}{ccc} X & \xrightarrow{\cong} & X \square_D D \\ \cong \downarrow & & \downarrow id_X \square_D \eta_D \\ C \square_C X & \xrightarrow{\delta_X \square_C id_X} & X \square_D h_D(X, D) \square_C X \end{array}$$

is commutative (Notice that $X \square_D (h_D(X, D) \square_C X) \cong (X \square_D h_D(X, D)) \square_C X$, since X_D is coflat). Hence $id_X \square_D \eta_D$ is an isomorphism, therefore, since X_D is faithfully coflat, η_D is an isomorphism.

Finally we show that $h_D(X, D) \square_C X$ is pure in $h_D(X, D) \otimes X$. Let $W \in R\text{-Mod}$ and consider the following commutative diagram, where μ_W is the canonical map of lemma 2.3

$$\begin{array}{ccccccc}
0 & \longrightarrow & X \square_D (h_D(X, D) \square_C (X \otimes W)) & \longrightarrow & X \square_D (h_D(X, D) \otimes X \otimes W) & \longrightarrow & X \square_D (h_D(X, D) \otimes C \otimes X \otimes W) \\
& & \uparrow id_X \square_C \mu_W & & \parallel & & \parallel \\
0 & \longrightarrow & X \square_D ((h_D(X, D) \square_C X) \otimes W) & \longrightarrow & X \square_D (h_D(X, D) \otimes X \otimes W) & \longrightarrow & X \square_D (h_D(X, D) \otimes C \otimes X \otimes W) \\
& & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
0 & \longrightarrow & (X \square_D h_D(X, D)) \square_C (X \otimes W) & \longrightarrow & (X \square_D h_D(X, D)) \otimes (X \otimes W) & \longrightarrow & (X \square_D h_D(X, D)) \otimes C \otimes X \otimes W,
\end{array}$$

in which the first and third rows are exact, since X_D is coflat. So the second row is also exact and hence $id_X \square_D \mu_W$ is an isomorphism. From lemma (2.3) it follows that $h_D(X, D) \square_C X$ is pure in $h_D(X, D) \otimes X$.

Now let $M \in \mathcal{M}^D$ and $N \in \mathcal{M}^C$, we have

$$\begin{aligned}
h_D(X, N \square_C X) &\cong (N \square_C X) \square_D h_D(X, D) \\
&\cong N \square_C (X \square_D h_D(X, D)) \quad (\text{since } h_D(X, D) \text{ is coflat in } {}^C\mathcal{M}) \\
&\cong N \square_C h_D(X, X) \\
&\cong N \square_C C \cong N, \quad \text{and} \\
h_D(X, M) \square_C X &\cong (M \square_D h_D(X, D)) \square_C X \\
&\cong M \square_D (h_D(X, D) \square_C X) \\
&\cong M \square_D D \cong M.
\end{aligned}$$

Therefore $-\square_C X$ is an equivalence with inverse $-\square_D h_D(X, -)$. ■

Definition 5.6. A C - D -bicomodule M is called invertible if the functor $-\square_C M : \mathcal{M}^C \rightarrow \mathcal{M}^D$ is an equivalence.

Corollary 5.7. Let C, D be two R -coalgebras. For a C - D -bicomodule X , the following are equivalent

- (1) X is invertible.
- (2) $X \square_D - : {}^D\mathcal{M} \rightarrow {}^C\mathcal{M}$ is an equivalence.
- (3) X_D is quasi-finite, faithfully coflat and an injector, and $e_D(X) \cong C$ as coalgebras.
- (4) ${}_C X$ is quasi-finite, faithfully coflat and an injector, and $e_C(X) \cong D$ as coalgebras.

In this case, C and D are Morita-Takeuchi equivalent.

6 Morita-Takeuchi Context

In this section we give a definition of Morita-Takeuchi context for coalgebras over rings. We also establish a correspondence between equivalences of comodule categories and strict Morita-Takeuchi contexts.

Definition 6.1. A *Morita-Takeuchi context* (D, C, M, N, f, g) consists of R -coalgebras D and C , bicomodules ${}_D M_C$, ${}_C N_D$, and bicomodule morphisms $f : D \longrightarrow M \square_C N$, $g : C \longrightarrow N \square_D M$, such that the following conditions hold

- (1) M and N are flat as R -modules.
- (2) $M \square_C N$ resp. $N \square_D M$ are pure in $M \otimes N$ resp. $N \otimes M$.
- (3) The diagrams

$$\begin{array}{ccc} M & \xrightarrow{\cong} & M \square_C C \\ \cong \downarrow & & \downarrow id \square_C g \\ D \square_D M & \xrightarrow{f \square_D id} & M \square_C N \square_D M, \end{array} \quad \begin{array}{ccc} N & \xrightarrow{\cong} & N \square_D D \\ \cong \downarrow & & \downarrow id \square_D f \\ C \square_C N & \xrightarrow{g \square_C id} & N \square_C M \square_D N. \end{array}$$

commute.

Remarks:. (a) Caenepeel [4] defined a Morita-Takeuchi context for coalgebras over rings which is the same as our definition but without the purity condition, with his definition it is not possible to prove that a strict Morita-Takeuchi context gives an equivalence between the involved coalgebras, since in his case the cotensor product is not associative.

(b) If R is a field, then conditions 1 and 2 are satisfied for all M and N . In this case our definition reduces to Takeuchi's one (see [14]).

(c) From condition 2 it follows that

$$\begin{aligned} X \square_C (N \square_D M) &\cong (X \square_C N) \square_D M, & Y \square_D (M \square_C N) &\cong (Y \square_D M) \square_C N, \\ N \square_D (M \square_C Z) &\cong (N \square_D M) \square_C Z, & M \square_C (N \square_D T) &\cong (M \square_C N) \square_D T, \end{aligned}$$

for all $X \in \mathcal{M}^C$, $Y \in \mathcal{M}^D$, $Z \in {}^C \mathcal{M}$ and $T \in {}^D \mathcal{M}$.

From a Morita-Takeuchi context (D, C, M, N, f, g) we have the following left exact functors

$$\begin{aligned} F &:= - \square_D M : \mathcal{M}^D \longrightarrow \mathcal{M}^C, \\ G &:= - \square_C N : \mathcal{M}^C \longrightarrow \mathcal{M}^D, \\ F' &:= M \square_C - : {}^C \mathcal{M} \longrightarrow {}^D \mathcal{M}, \\ G' &:= N \square_D - : {}^D \mathcal{M} \longrightarrow {}^C \mathcal{M}, \end{aligned}$$

and the following natural transformations

$$\begin{aligned}\lambda : id_{\mathcal{M}^C} &\rightarrow FG, & \zeta : id_{\mathcal{M}^D} &\rightarrow GF, \\ \psi : id_{D\mathcal{M}} &\rightarrow F'G' & \delta : id_{C\mathcal{M}} &\rightarrow G'F',\end{aligned}$$

which are defined as follows

$$\begin{aligned}\lambda_X &: X \cong X \square_C C \xrightarrow{id \square_C g} X \square_C (N \square_D M) \cong (X \square_C N) \square_D M = FG(X), \\ \zeta_Y &: Y \cong Y \square_D D \xrightarrow{id \square_D f} Y \square_D (M \square_C N) \cong (Y \square_D M) \square_C N = GF(Y), \\ \psi_T &: T \cong D \square_D T \xrightarrow{f \square_D id} (M \square_C N) \square_D T \cong M \square_C (N \square_D T) = F'G'(T), \\ \delta_Z &: Z \cong C \square_C Z \xrightarrow{g \square_C id} (N \square_D M) \square_C Z \cong N \square_D (M \square_C Z) = G'F'(Z),\end{aligned}$$

Lemma 6.2. *Let (D, C, M, N, f, g) be a Morita-Takeuchi context.*

(1) *If f is bijective, then*

- (a) $G = -\square_C N : \mathcal{M}^C \longrightarrow \mathcal{M}^D$ is left adjoint to $F = -\square_D M : \mathcal{M}^D \longrightarrow \mathcal{M}^C$.
- (b) $F' = M \square_C - : {}^C\mathcal{M} \longrightarrow {}^D\mathcal{M}$ is left adjoint to $G' = N \square_D - : {}^D\mathcal{M} \longrightarrow {}^C\mathcal{M}$.

(2) *If g is bijective, then*

- (a) $F = -\square_D M : \mathcal{M}^D \longrightarrow \mathcal{M}^C$ is left adjoint to $G = -\square_C N : \mathcal{M}^C \longrightarrow \mathcal{M}^D$.
- (b) $G' = N \square_D - : {}^D\mathcal{M} \longrightarrow {}^C\mathcal{M}$ is left adjoint to $F' = M \square_C - : {}^C\mathcal{M} \longrightarrow {}^D\mathcal{M}$.

(3) *If f and g are bijective, then*

- (a) $G = -\square_C N$ is an equivalence with inverse $F = -\square_D M$.
- (b) $G' = N \square_D -$ is an equivalence with inverse $F' = M \square_C -$.

In this case C and D are Morita-Takeuchi equivalent coalgebras.

Proof: (1) Assume f is bijective

- (a) We have the natural transformation $\lambda : id_{\mathcal{M}^C} \longrightarrow FG$. Since f is bijective $\zeta^{-1} : GF \longrightarrow id_{\mathcal{M}^D}$ is a natural transformation. We are done if we show that each of the composition of morphisms

$$\begin{aligned}X \square_C N &\cong X \square_C C \square_C N \xrightarrow{id \square_C g \square_C id} X \square_C N \square_D M \square_C N \xrightarrow{id \square_C id \square_D f^{-1}} X \square_C N, \\ Y \square_D M &\cong Y \square_D M \square_C C \xrightarrow{id \square_D id \square_C g} Y \square_D M \square_C N \square_D M \xrightarrow{id \square_D f^{-1} \square_D id} Y \square_D M,\end{aligned}$$

gives the identity on $X \square_C N$ resp. $Y \square_D M$, for all $X \in \mathcal{M}^C$ and $Y \in \mathcal{M}^D$. We have

$$\begin{aligned}
(id \square_C id \square_D f^{-1})(id \square_C g \square_C id) &= id \square_C (id \square_D f^{-1})(g \square_C id) \\
&= id \square_C (id \square_D f^{-1})(id \square_D f) \\
&= id \square_C (id \square_D id) \\
&= id_{X \square_C N}.
\end{aligned}$$

The other composition is similar.

(b) is similar to (a).

(2) Similar to (1)

(3) Assume f and g are bijective, let $X \in \mathcal{M}^C$, $Y \in \mathcal{M}^D$. We have

$$\begin{aligned}
FG(X) &= (X \square_C N) \square_D M \\
&\cong X \square_C (N \square_D M) \\
&\cong X \square_C C \cong X, \quad \text{und} \\
GF(Y) &= (Y \square_D M) \square_C N \\
&\cong Y \square_D (M \square_C N) \\
&\cong Y \square_D D \\
&\cong Y.
\end{aligned}$$

Hence F is a category equivalence with inverse G . Similarly one can show that G' is an equivalence with inverse F' . ■

Corollary 6.3. Let (D, C, M, N, f, g) be a Morita-Takeuchi context.

(1) If f is bijective, then

- (a) The comodules M_C and ${}_C N$ are quasi-finite, coflat and injectors.
- (b) $e_C(M) \cong D$ and $e_C(N) \cong D$ as coalgebras, ($e_C(M)$ is the coendomorphism coalgebra of M).
- (c) g is bijective iff ${}_C N$ is faithfully coflat iff M_C is faithfully coflat.

(2) If g is bijective, then

- (a) The comodules N_D and ${}_D M$ are quasi-finite, coflat and injectors.
- (b) $e_D(N) \cong C$ and $e_D(M) \cong C$ as coalgebras.
- (c) f is bijective iff ${}_D M$ is faithfully coflat iff N_D is faithfully coflat.

(3) If f and g are bijective, then

- (a) The comodules M_C , N_D , ${}_C N$ and ${}_D M$ are quasi-finite, faithfully coflat and injectors.
- (b) $e_C(M) \cong D$, $e_C(N) \cong D$, $e_D(N) \cong C$, and $e_D(M) \cong C$ as coalgebras.

Proof: (1) Assume that f is bijective.

- (a) From the fact that $G = -\square_C N$ is left adjoint to $F = -\square_D M$, it follows that M_C is quasi-finite and that G is right exact. Since C and N are flat R -modules we know that G is also left exact (see 2.2). So ${}_C N$ is coflat. Similarly one can show that M_C is coflat and that ${}_C N$ is quasi-finite. The assertion that these comodules are injectors follows from the fact that a functor between abelian categories which has an exact left adjoint preserves injective objects.
- (b) Follows from the construction of the coendomorphism coalgebra.
- (c) If g is bijective, then $-\square_C N$ is an equivalence, hence ${}_C N$ is faithfully coflat. If ${}_C N$ is faithfully coflat, then $g\square_C id_N : C\square_C N \longrightarrow (N\square_D M)\square_C N$ is an isomorphism, hence g is an isomorphism. The other assertion is similar.

(2) Similar to (1)

(3) Follows directly from (1) and (2). ■

Definition 6.4. A Morita-Takeuchi context (D, C, M, N, f, g) in which f and g are both bijective is called a *strict Morita-Takeuchi context*.

In lemma 6.2 we have seen that the involved coalgebras in a strict Morita-Takeuchi context are Morita-Takeuchi equivalent. The converse is also true as the following theorem shows

Theorem 6.5. *For two coalgebras C , D , the following are equivalent*

- (1) C is Morita-Takeuchi equivalent to D .
- (2) There exists a strict Morita-Takeuchi context (D, C, M, N, f, g)

Proof: We have only to show (1) $\Rightarrow$ (2). Let $F : \mathcal{M}^C \rightarrow \mathcal{M}^D$ be an equivalence with inverse $G : \mathcal{M}^D \rightarrow \mathcal{M}^C$, and let $\phi : id_{\mathcal{M}^C} \longrightarrow GF$, $\psi : id_{\mathcal{M}^D} \longrightarrow FG$ be the natural transformation belonging to this equivalence. From 4.6 we have $F \cong -\square_C F(C)$ and $G \cong -\square_D G(D)$. Let $f := \phi_C : C \longrightarrow F(C)\square_D G(D)$, and $g := \psi_D : D \longrightarrow G(D)\square_C F(C)$.

Claim, $(D, C, G(D), F(C), f, g)$ is a strict Morita-Takeuchi context. The purity and flatness conditions follow from the general properties of the bicomodules $F(C)$ and $G(D)$. From the definition of f, g and that the comultiplication of C and D are bilinear maps it follows that f and g are bilinear.

The commutativity of the diagrams follows from the relations

$$F\phi = \psi H, \quad \text{and} \quad G\psi = \phi G.$$

So we have a strict Morita-Takeuchi context. ■

Let X be a right C -comodule, and assume that it is quasi-finite, faithfully coflat and an injector. We will construct a strict Morita-Takeuchi context derived from X . In 3 we have seen that X is a $e_C(X)$ - C -bicomodule and that $h_D(X, C)$ is a C - $e_C(X)$ -bicomodule, where $h_D(X, -)$ is the cohom functor. Now let f be the D -colinear map (see 3.5) $f := \delta_X : h_D(X, X) \longrightarrow X \square_C h_D(X, C)$ and $g := \eta_C : C \longrightarrow h_D(X, C) \square_{e_C(X)} X$ (see 3.7).

Proposition 6.6. *Let X be as above, then $(e_C(X), C, {}_{e_C(X)}X_C, {}_Ch_D(X, C)_{e_C(X)}, f, g)$ is a strict Morita-Takeuchi context.*

Proof: It is clear that X is flat in R -Mod. From 3.11 it follows that $h_D(X, C)$ is also a flat R -module, and that $f = \delta_X$ is bijective. As in the proof of 5.5 it is easy to show that g is bijective and that $h_D(X, C) \square_{e_C(X)} X$ is pure in $h_D(X, C) \otimes X$. Since $h_D(X, -)$ is exact, $X \square_C h_D(X, C)$ is pure in $X \otimes h_D(X, C)$. The commutativity of the diagrams follows from the defining properties of f and g . ■

Corollary 6.7. *Let X be as above, then C is Morita-Takeuchi equivalent to $e_C(X)$, the coendomorphism coalgebra of X .*

7 Coalgebras over QF-Rings

In this section we consider comodule categories for coalgebras over QF-rings. We will show that Takeuchi's description of the cohom functor is valid in this case under the condition that the coalgebra is projective over the ground ring. From now on we assume that all coalgebras are projective over R .

We need the following two lemmas whose proofs can be found in [16].

Lemma 7.1. *Let D be an R -coalgebra, M a D -comodule. Then*

- (1) *every finite subset of M is contained in a subcomodule of M which is finitely generated as R -module.*
- (2) *The following are equivalent*
 - (a) *D is finitely generated as R -module.*
 - (b) $\mathcal{M}^D = D^* \text{-Mod} (= \sigma[D^*D]).$

Let R be a noetherian ring, D an R -coalgebra. From lemma 7.1, it follows that every D -comodule is the direct limit of its subcomodules, that are finitely presented as R -modules. From this and the com-cotensor relations (see [2]), one obtains some relations between injectivity and coflatness for coalgebras over QF-rings.

Lemma 7.2. *Let D be a coalgebra over a QF-ring. A D -comodule M , which is flat as R -module is injective (resp. an injective cogenerator) if and only if it is coflat (resp. faithfully coflat)*

From this lemma and 1.2 it follows that over a QF-ring the coalgebra itself is an injective cogenerator as a right and left comodule.

Corollary 7.3. Let D be a coalgebra over a QF-ring, M a right D -comodule, which is flat as R -module. If M is injective in $\mathcal{M}^D$, then it is an injector.

Proof: Let M be an injective D -comodule, W an injective R -module. Since R is QF, W is flat and M is coflat. Now

$$(W \otimes M) \square_D - \cong W \otimes (M \square_D -) : {}^D\mathcal{M} \rightarrow R\text{-Mod}.$$

So the functor $(W \otimes X) \square_D -$ is exact, i.e. $W \otimes X$ is coflat, hence it is injective. ■

Proposition 7.4. *Let D be a coalgebra over a coherent ring, X a quasi-finite D -comodule, P a D -comodule. If P is finitely presented as R -module, then $\text{Com}_D(P, X)$ is finitely presented.*

Proof: For every index set Λ , R^Λ is flat. Consider

$$\begin{aligned} R^\Lambda \otimes \text{Com}_D(P, X) &\cong \text{Com}_D(P, R^\Lambda \otimes X) \\ &\cong \text{Hom}_R(h_D(X, P), R^\Lambda) \\ &\cong \text{Hom}_R(h_D(X, P), R)^\Lambda \\ &\cong \text{Com}_D(P, R \otimes X)^\Lambda \\ &\cong \text{Com}_D(P, X)^\Lambda. \end{aligned}$$

the composition of these isomorphisms is the canonical map

$$R^\Lambda \otimes \text{Com}_D(P, X) \rightarrow \text{Com}_D(P, X)^\Lambda, (r_i) \otimes f \mapsto (r_i f)$$

Hence $\text{Com}_D(P, X)$ is finitely presented as R -module. ■

Now we prove our main result in this section, which gives Takeuchi's representation of the cohom functor for coalgebras over rings.

Theorem 7.5. *Let C, D be coalgebras over a QF-ring R , X a C - D -bicomodule. If X_D is quasi-finite, injective cogenerator and $e_D(X) \cong C$ as coalgebras, then the functor*

$$-\square_C X : \mathcal{M}^C \rightarrow \mathcal{M}^D$$

is an equivalence with inverse

$$h_D(X, -) \cong -\square_D h_D(X, D) : \mathcal{M}^D \rightarrow \mathcal{M}^C$$

Moreover, for every $M \in \mathcal{M}^D$, the cohom functor is given by

$$h_D(X, M) = \lim_{\substack{\longrightarrow \\ \Lambda}} \text{Com}_D(M_\lambda, X)^*,$$

where $\{M_\lambda\}_\Lambda$ is the family of subcomodules of M that are finitely presented as R -modules.

Proof: From lemma 7.2 and corollary 5.7 it follows that $-\square_C X : \mathcal{M}^C \rightarrow \mathcal{M}^D$ is an equivalence with inverse $h_D(X, -) \cong -\square_D h_D(X, D) : \mathcal{M}^D \rightarrow \mathcal{M}^C$. For the other assertion consider the following commutative diagram of categories and functors

$$\begin{array}{ccc} \mathcal{M}_f^D & \begin{array}{c} \xrightarrow{\text{Com}_D(-, X)} \\ \xleftarrow{\text{Com}_C(-, X)} \end{array} & {}^C\mathcal{M}_f \\ & \searrow \begin{array}{c} -\square_C X \\ h_D(X, -) \end{array} & \downarrow \begin{array}{c} (-)^* \\ (-)^* \end{array} \\ & & \mathcal{M}_f^C, \end{array}$$

where $\mathcal{M}_f^C, \mathcal{M}_f^D$ and ${}^C\mathcal{M}_f$ are the categories of comodules that are finitely presented as R -modules.

Let $P \in \mathcal{M}_f^D$, then we have

$$\begin{aligned} \text{Com}_D(P, X)^* \square_C X &\cong \text{Com}_C(\text{Com}_D(P, X), X) \\ &\cong \text{Com}_C(X \square_D P^*, X) \\ &\cong \text{Com}_D(P^*, h_D(X, D) \square_C X) \\ &\cong \text{Com}_D(P^*, D) \cong P, \end{aligned}$$

here we have used the facts that $\text{Com}_D(P, X)$ is finitely presented, and that $h_D(X, D) \square_C -$ is right adjoint to $X \square_D -$.

For $Q \in \mathcal{M}_f^C$ we have

$$\begin{aligned} \text{Com}_D(Q \square_C X, X)^* &\cong \text{Com}_C(Q, h_D(X, X))^* \\ &\cong \text{Com}_C(Q, C)^* \\ &\cong Q^{**} \cong Q, \end{aligned}$$

here we used the fact that $-\square_C X$ is left adjoint to $h_D(X, -)$. Hence the functor $\text{Com}_D(-, X)^* : \mathcal{M}_f^D \rightarrow \mathcal{M}_f^C$ is left adjoint to the functor $-\square_C X : \mathcal{M}_f^C \rightarrow \mathcal{M}_f^D$ and therefore is isomorphic to $h_D(X, -)$.

Now let $M \in \mathcal{M}^D$, so $M = \varinjlim_{\Lambda} M_{\lambda}$, where $\{M_{\lambda}\}_{\Lambda}$ is the family of subcomodules of M that are finitely presented as R -modules. We have

$$\begin{aligned} h_D(X, M) &\cong h_D(X, \varinjlim_{\Lambda} M_{\lambda}) \\ &\cong \varinjlim_{\Lambda} h_D(X, M_{\lambda}) \\ &\cong \varinjlim_{\Lambda} \text{Com}_C(M_{\lambda}, X)^*. \end{aligned}$$

■

Summarizing the results of this section together we get a characterization for the equivalences of comodule categories over QF-rings that agrees with Takeuchi's results.

Corollary 7.6. Let C, D be coalgebras over a QF-ring R , X a C - D -bicomodule. The following are equivalent

1. X is invertible.
2. The functor $X \square_D - : {}^D\mathcal{M} \rightarrow {}^C\mathcal{M}$ is an equivalence.
3. X_D is quasi-finite, injective cogenerator and $e_D(X) \cong C$ as coalgebras.
4. ${}_C X$ is quasi-finite, injective cogenerator and $e_C(X) \cong D$ as coalgebras.

Moreover, for every $M \in \mathcal{M}^D$, the cohom functor is given by

$$h_D(X, M) = \varinjlim_{\Lambda} \text{Com}_D(M_{\lambda}, X)^* \cong \varinjlim_{\Lambda} (X \square_D M_{\lambda}^*)^*,$$

where $\{M_{\lambda}\}_{\Lambda}$ is the family of subcomodules of M that are finitely presented as R -modules.

Acknowledgement

This article is part of my Ph.D thesis at Düsseldorf Universität, Germany under the supervision of Prof. Dr. Robert Wisbauer, many thanks to him for his continuous support and encouragement. I would also thank Prof. José Gómez-Torrecillas for the fruitful discussions.

References

- [1] K. Al-Takhman. *Äquivalenzen zwischen Komodulkategorien von Koalgebren Über Ringen*. Ph. D. Dissertation, H.H.U. Düsseldorf, Germany, 1999.
- [2] K. Al-Takhman. The com and cotensor functors for coalgebras over rings. *proceedings of the third international palestinian conference on mathematics and mathematics education, Bethlehem, Palestine*, August 2000.
- [3] F. Anderson and K. Fuller. *Rings and Categories of Modules*. Springer-Verlag, Berlin, 1973.
- [4] S. Caenepeel. *Brauer Groups, Hopf algebras and Galois theory*, volume 4 of *K-Monographs in Mathematics*. Kluwer Academic Publishers, 1998.
- [5] S. Dăscălescu, C. Năstăsescu, S. Raianu, and F. Van Oystaeyen. Graded coalgebras and Morita-Takeuchi contexts. *Tsukuba J. Math.*, 19(2):395–407, 1995.
- [6] L. Grunenfelder and R. Paré. Families parametrized by coalgebras. *J. Algebra*, 107:316–375, 1987.
- [7] F. Guzman. *Cointegration and Relative Cohomology for Comodules*. Ph. D. Dissertation, Syracuse University, 1985.
- [8] F. Guzman. Cointegration, relative cohomology for comodules, and coseparable corings. *J. Algebra*, 126:211–224, 1989.
- [9] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Ann. Math.*, 81:211–264, 1965.
- [10] Popesco. *Abelian categories with applications to rings and modules*. Academic Press, London, 1973.
- [11] H.-J. Schneider. Principal homogeneous spaces for arbitrary Hopf algebras. *Israel J. Math.*, 72(1-2):167–195, 1990.
- [12] B. Stenström. *Rings and modules of quotients*, volume No. 237 of *Lecture Notes in Mathematics*. Springer Verlag, 1971.
- [13] M. Takeuchi. Formal schemes over fields. *Comm. Algebra*, 5:1483–1528, 1977.
- [14] M. Takeuchi. Morita theorems for categories of comodules. *J. Fac. Sci. Univ. Tokyo*, 24:629–644, 1977.

- [15] R. Wisbauer. *Modules and Algebras: Bimodule Structure and Group Actions on Algebras*. Pitman Mono. PAM 81, Addison Wesley Longman, Essex, 1996.
- [16] R. Wisbauer. Introduction to Coalgebras and Comodules. Lecture Notes, August 1998.